

BUPACIN

Buparvaquone Injection 50 mg/ml

COMPOSITION :

Each ml contains :

Buparvaquone	50 mg
Sorbitan mono oleate B.P	100 mg
Oil Base	q.s.

INDICATION :

It is indicated for theileriosis (East Coast Fever, Corridor Disease, Zimbabwean Theileriosis, Tropical Theileriosis) caused by intracellular protozoan parasites *Theileria parva*, *Theileria parvabovis*, *Theileria parva lawrencei*, *T. annulata*, *T. mutans* and *T. orientalis sergenti* in Cattle. It is active against both the schizont and piroplasm stage of *Theileria* spp., and may be used during the incubation period of the disease or when clinical signs are apparent.

DOSAGE AND ADMINISTRATION :

A single dose of 1 ml BUPACIN Injection per 20 kg bodyweight. (2.5 mg buparvaquone per kg bodyweight). In severe cases the treatment may be repeated within 48-72 hours. Not more than 10 ml should be injected into a single site. Successive Injection should be administered at different sites.

ADMINISTRATION :

Intramuscular Injection into the neck muscles.

CONTRAINDICATION :

Due to the inhibiting effects of theileriosis on the immune system, Vaccination should be delayed until the animal has recovered from theileriosis.

WITHDRAWAL PERIOD :

Milk : 2 days
Meat : 42 days



Intracin

Pharmaceutical Pvt. Ltd.

C-1, B-53, G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

