

# BUTACIN

## Butaphosphan and Cyanocobalamin Injection

### COMPOSITION :

#### Each ml contains :

Butaphosphan	100 mg
Cyanocobalamin B.P	50 mcg
Methyl Hydroxy Benzoate B.P	0.1% w/w
Water for Injections B.P	q.s.

### INDICATION :

BUTACIN is indicated for debilitation by acute or chronic metabolism disorders that result from poor nutrition, inadequate management or disease (e.g. developmental and nutritional disorders in young animals due to rearing disease, and (secondary) ketosis in cows). It can be used for metaphylaxis of infertility, puerperal diseases and in support of sterility treatment. It acts as a roborant in cases of stress, overexertion, exhaustion and reduced resistance and as a tonic in cases of weakness, secondary anaemia and chilling. BUTACIN additionally supports muscular physiology, the treatment of infertility, and tetany and paresis as an adjunct to calcium and magnesium therapy.

### DOSAGE AND ADMINISTRATION :

In the form of Intravenous, Intramuscular and subcutaneous Injection:

Horses and cattle	: 10.0 - 25.0 ml.
Foals and calves	: 5.0 - 12.0 ml.
Sheep and goats	: 2.5 - 8.0 ml.
Lambs	: 1.5 - 2.5 ml.
Pigs	: 2.5 - 10.0 ml.
Gilts	: 1.0 - 2.5 ml.
Dogs	: 0.5 - 5.0 ml.

Cats, fur-bearing animals : 0.5 - 2.5 ml

If necessary, repeat the treatment next day.

For birds Butacin added to the drinking water at the rate of:

Chicks and young 1.0 - 1.5 ml to 1 liter

Broilers and laying hens 2.0 - 3.0 ml to 1 liter and fed within 4 - 5 days.

### WITHDRAWAL PERIOD :

Meat : 0 days

Milk : 0 days

### STORAGE :

Store at a temperature not exceeding 30°C.

Protect from sunlight.

IPPL

## Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,  
NADIAD-387001 (GUJARAT) INDIA.  
[www.intracin.com](http://www.intracin.com)

