# CIT-C

# Ascorbic Acid Injection 500 mg/ml

#### **COMPOSITION:**

Each ml contains:

Ascorbic acid (Vitamin C)

Methyl hydroxybenzoate B.P

Propyl hydroxybenzoate B.P

Water for Injection B.P

200.0 mg
0.8 mg
0.2 mg
q.s.

#### **INDICATIONS:**

In Cattle, Sheep, Goats, Horses, Pigs, Dogs, Cats and Poultry

- Treatment of vitamin C deficiency.
- Adjuvant therapy for asthenia, notably in cases of infection or stress.

# **CONTRA-INDICATIONS:**

Do not use in case of hypersensitivity to the active substances or to any of the excipients.

#### **DOSAGE AND ADMINISTRATION:**

IV. IM and oral.

Daily dose: 20 to 60 mg ascorbic acid per kg body weight per day, corresponding to:

HORSES - CATTLE : Adults from 50 to 100 ml Youngs from 10 to 30 ml

SHEEP, GOATS, PIGS from 5 to 25 ml

Oral administration is preferable for chronic conditions or as an addition to parenteral administration. Orally, the product can be administered on its own, mixed uniformly into feed, or diluted in water (do not mix with milk).

### **WITHDRAWAL PERIOD:**

Cattle, Sheep, Goats, Horses : Meat and offal : zero days. Milk : zero days.

Pigs:

Meat and offal: zero days.

Poultry:

Meat and offal : zero days. Eggs : zero days

# **WARNING:**

Special warnings for each target species: None Special Precautions for use in animals: None

Special precautions to be taken by the person administering the veterinary mdicinal product

to aniamals : None Other precautions : None

Use during pregnancy, lactation or lay: In the absence of any teratogenesis studies for animals, the use of vitamin C is not recommended during gestation unless absolutly necessary. Interaction with other medicinal products and other forms of

Interaction : None Overdose : Not known.

#### STORAGE:

Do not store above 30°C. Protect from light. Shelf life after first opening the container : 28 days.

Keep out of reach and sight of children.



C-1,B-53,G.I.D.C. ESTATE, NADIAD-387001 (GUJARAT) INDIA.

