

CRAMINE

Chlorphenamine Injection B.P 10 mg/ml

COMPOSITION :

Each ml contains :

Chlorphenamine maleate B.P	10 mg
Phenyl mercuric nitrate B.P (As Preservative)	0.002 % w/v
Water for Injections B.P	q.s.

INDICATION :

1. Allergic reactions and allergic disorders with manifestation of acute respiratory signs.
2. Rhinitis, dyspnoea, pulmonary emphysema, pulmonary oedema, asthma and hay fever.
3. Eczema, dermatitis, urticaria, insect bite, tail eczema in Horses, toxic hoof corn
4. Ruminal atony, tympany and bloat in ruminant.
5. Puerperal toxæmia and secondary placental retention, acute septic metritis and gangrenous mastitis.
6. Reactions and other drug allergies and anaphylactic shock.

CONTRAINDICATION :

Do not use in animals known to be hypersensitive to chlorphenamine.
Do not use in renal and hepatic insufficiency.

DOSAGE AND ADMINISTRATION :

Cattle & Horse : 5-10 ml

Sheep & Goat : 0.5 - 2 ml. By Intramuscular route.

Canine : 0.5 - 2 ml

Poultry : 0.5 - 1 ml. By Intramuscular route or as directed by veterinarian.

WITHDRAWAL PERIOD :

Nil

STORAGE :

Store at a temperature not exceeding 30°C.
Protect from light.

Veterinary Medicines - Injectables

IPPL

Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

