



# DEXACIN-F

## Dexamethasone with Furosemide Injection

### COMPOSITION :

#### Each ml contains :

Furosemide	B.P	20 mg
Dexamethasone Sodium Phosphate	B.P	0.63 mg
Phenol	B.P	0.5% v/v
Water for Injection	B.P	q.s.

### INDICATIONS :

For treatment of edemas of different etiology in Bovines, Sheep and Horses.

Especially indicated for treatment of post partum udder edema in Cows.

### SPECIAL PRECAUTIONS AND CONTRA-INDICATIONS :

Concomitant use of aminoglycoside antibiotics should be avoided due to possible potentiation of nephrotoxic effects of furosemide.

Do not exceed recommended dose and duration of treatment.

As with all diuretic drugs, over dosage for prolonged periods of time may cause electrolyte imbalance.

In this case, treatment should be suspended and electrolyte balance restored.

Do not administer to animals with compromised liver or kidney function.

Do not administer to horses used for human consumption.

### DOSAGE AND ADMINISTRATION :

Administer by intramuscular or subcutaneous route, at 0.5 ml per 20 kg of body weight. Dose can be repeated after 24 hr.

Duration of treatment may vary according to each clinical case; usually 2 doses with a 24 hr interval are sufficient.

### WITHDRAWAL PERIOD :

Meat & Offals : 3 days

Milk : 2 days

### STORAGE :

Store at a temperature not exceeding 30°C.

Protect from direct sunlight.

Veterinary Medicines - Injectables



## Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,  
NADIAD-387001 (GUJARAT) INDIA.

[www.intracin.com](http://www.intracin.com)

