

DEXAZIDE

Dexamethasone with Hydrochlorothiazide Injection



COMPOSITION :

Each ml contains :

Dexamethasone sodium phosphate	B.P	0.50 mg
Hydrochlorothiazide	B.P	50.00 mg
Propylene glycol	B.P	0.15 ml
Benzyl alcohol	B.P	0.01 ml
Water for Injection	B.P	q.s.

INDICATIONS :

CATTLE :

- Congestion and oedema of the udder.
- Early persistent lactation edema.
- Pulmonary congestion and edema.
- Edematous infiltration of surgical wounds.
- Edema in allergic conditions.

HORSES :

- Generalised congestion and oedema
- Oedema of sheath Anasarca
- Oedema in allergic conditions

DOSAGE AND ADMINISTRATION :

Dexazide injection may be administered by Intravenous, intramuscular, subcutaneous injection.

Cattle and adult Horses :

Preventative treatment : 10 ml daily for 3 days.

Curative treatment :

Congestion and mild oedema : 10 ml daily for 2 or 3 days

Congestion and severe oedema : 20 ml daily for 2 days and 10 ml the third day.

Foals - Calves : 2 ml per 40 - 50 kg bodyweight daily for 3 days

WITHDRAWAL PERIOD :

Cattle : Meat and offal : 28 days

Milk : 7 days

Treated Horses may never be slaughtered for human consumption.

STORAGE :

Store at a temperature not exceeding 30°C.

Protect from direct sunlight.



Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

