

FLASIN WS

Tylosin Tartrate Powder for oral administration



COMPOSITION :

Each gram contains :

Tylosin tartrate B.P 1000 mg

DESCRIPTION :

Tylosin is a macrolide antibiotic with a bacteriostatic action against Gram-positive and Gram-negative bacteria like Campylobacter, Mycoplasma, Pasteurella, Staphylococcus, Streptococcus and Treponema spp. and Mycoplasma.

INDICATIONS :

Gastrointestinal and respiratory infections caused by tylosin sensitive micro-organisms, like Campylobacter, Mycoplasma, Pasteurella, Staphylococcus, Streptococcus and Treponema spp. in Calves, Goats, Poultry, Sheep and Swine.

CONTRA-INDICATIONS :

Hypersensitivity to tylosin.
Concurrent administration of penicillins, cephalosporins, quinolones and cycloserine.
Administration to animals with an active microbial digestion.

SIDE EFFECTS :

Diarrhoea, epigastric pain and skin sensitization can occur.

DOSAGE AND ADMINISTRATION :

For oral administration:

Calves, Goats and Sheep : Twice daily, 5 g per 220 - 250 kg body weight for
5 - 7 days.

Poultry : 1 kg per 1500 - 2000 litres of drinking water for
3 - 5 days.

Swine : 1 kg per 3000 - 4000 litres of drinking water for
5 - 7 days.

Note : for pre-ruminant Calves, Lambs and Kids only.

WITHDRAWAL PERIOD :

For meat :

Calves, Goats, Poultry and Sheep : 5 days.

Swine : 3 days.



Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

