

FUTIC

Furosemide Injection B.P 5% w/v

COMPOSITION :

Each ml contains :

Furosemide Sodium B.P
equi.to Furosemide 50 mg
Benzyl Alcohol B.P 1% v/v
(as preservative)
Water for Injection B.P q.s.

INDICATIONS :

It is used for treatment of congestive cardiomyopathy, pulmonary edema, hypercalcuric nephropathy, uremia and adjunctive therapy in hyperkalemia & occasionally as antihypertensive agent. In Cattle it is used for the treatment of post parturient udder edema & associated structures. In race Horses, it helps to prevent or to reduce epistaxis.(exercise included pulmonary hemorrhage).

CONTRA-INDICATIONS :

Do not use the product in cases of acute glomerular nephritis, renal failure with anuria, electrolyte deficiency disease or overdosage with digitalis. Do not use concurrently with aminoglycoside antibiotic treatment.

DOSAGE AND ADMINISTRATION :

Dogs and Cat : 1 to 2 mg/kg bodyweight once or twice daily at 6 to 8 hour intervals.
Cattle : 0.5 - 1 mg/kg bodyweight.
Horses : 0.5 - 1 mg/kg bodyweight once or twice daily at 6 to 8 hour intervals.

Administration: Intramuscular or Intravenous route

WITHDRAWAL PERIOD :

Do not use in Horses intended for human consumption. Treated Horses may never be slaughtered for human consumption.
Cattle : Meat & Milk : 48 hours.

STORAGE :

Store at a temperature not exceeding 30°C.
Protect from light.



Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

