

IMIX

Imidocarb Dipropionate Aqueous Injection Solution

COMPOSITION :

Each ml contains :

Imidocarb Dipropionate	120 mg
Benzyl Alcohol B.P	2.0% w/v
Water For Injections B.P	q.s.

INDICATIONS :

IMIX is used for the treatment of babesiosis in Cattle, Sheep, Horses and Dogs and of anaplasmosis in Cattle. IMIX will also treat ehrlichiosis in Dogs. For Subcutaneous or Intramuscular Injection.

DOSAGE AND ADMINISTRATION :

Dose according to body weight:

Cattle : 1 ml per 100 kg bodyweight. In treating mixed infections due to Anaplasma and Babesia, administer 2.5 ml per 100 kg body weight.

Sheep : 0.5 ml per 50 kg bodyweight.

Horses : 2 ml per 100 kg bodyweight. (Intramuscular injection is preferred). In most cases, a single dose will effect a complete cure, but treatment of B. equi infections in Horses may require 2 doses at an interval of 24 hours.

Dogs : 0.25 ml per 10 kg body weight. In treating mixed infection due to Ehrlichina and Babesia, administer 0.5 ml per 10 kg bodyweight, 2 doses at an interval of 14 days.

It is advised to screen the blood after one treatment for presence of the parasite. Repeat treatment for presence of the parasite. Repeat treatment if the test is positive.

WITHDRAWAL PERIOD :

Meat : 28 days

Milk : 7 days.

STORAGE :

Store at a temperature not exceeding 25°C.

Protect from light.



Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.

www.intracin.com

