

IMOLY 20% LA

Oxytetracycline Injection Long Acting 20% w/v

COMPOSITION :

Each ml contains :

Oxytetracycline Dihydrate B.P
eq. to Anhydrous Oxytetracycline 200 mg
in 2-Pyrrolidone Vehicle system q.s.

INDICATIONS :

IMOLY 20% is indicated in the treatment and control of disease in Cattle, Sheep and Pigs caused by or associated with organisms sensitive to Oxytetracycline. For the treatment and control of pasteurellosis and pneumonia caused by oxytetracycline sensitive organisms and as an aid in the treatment of infectious bovine keratoconjunctivitis (New Forest Disease) due to oxytetracycline sensitive strains of *Moraxella bovis*. This product may also be of value for foul-in-the-foot.

CONTRA-INDICATIONS :

Not recommended for Cats, Dogs, Horses and Donkeys.

The use of this product during the period of tooth development including late pregnancy may lead to tooth discoloration.

DOSAGE AND ADMINISTRATION :

For Intramuscular or Subcutaneous administration:

General : 1 ml per 10 kg body weight.

This dosage can be repeated after 48 hours when necessary.

Do not administer more than 20 ml in Cattle, more than 10 ml in Swine and more than 5 ml in Calves, Goats and Sheep per Injection site.

WITHDRAWAL PERIOD :

Milk for human consumption should not be taken during treatment. Milk intended for human consumption may be taken only after 7 days from the last treatment. Not for use in ewes producing milk for human consumption. Animals may only be slaughtered for human consumption after last treatment, according to the below table.

| Species | Meat withdrawal periods |
|----------------|-------------------------|
| Cattle & Sheep | 28 days |
| Pigs | 35 days |

STORAGE :

Store at a temperature not exceeding 30°C.

Protect from light.

IPPL

Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

