

IROFLOX SA

Enrofloxacin Injection 10% w/v (Vet.)

COMPOSITION:

Each ml contains:

Enrofloxacin USP 100 mg
Benzyl Alcohol B.P 1.5% v/v
[as preservative]
Water for Injection B.P q.s.

INDICATION:

Shipping fever pneumonia (H.S), Pneumonia, Pleuropneumonia, Pseudorinderpest and Urinary infections, mastitis caused by mycoplasma spp. Respiratory and enteric diseases (Pasteurellosis, Mycoplasmosis, Colibacillosis, Colisepticaemia and secondary bacterial complication during viral infections).

Mastitis caused by susceptible microorganisms and renal infection.

IROFLOX SA is effective against the following microorganism : Mycoplasma spp., E.coli, Salmonella spp., Haemophilus spp., Pasteurella spp., Klebsiella spp., Proteus spp., Vibrio parahaemolyticus and Staphylococcus aureus.

DOSAGE:

Cattle, Buffaloes, Sheep, Goats and Camels:

As directed by the veterinarian. OR 1 ml per 20-40 kg body weight depending upon the severity of infection. (2.5-5 mg of Enrofloxacin 1kg body weight) the second administration should be administered after 72 hours, if required.

ROUTE:

For I.M./SC/Slow I.V. Use.

WITHDRAWAL PERIOD:

Animals must not be slaughtered and milk should not be used for human consumption during treatment.

The withdrawal periods after the last treatment are:

Cattle:

For Intravenous route : Meat : 4 days
Milk : 72 hours

For Subcutaneous route : Meat : 14 days
Milk : 84 hours

STORAGE :

Store at a temperature not exceeding 30°C.
Protect from light.

IPPL

Intracin

Pharmaceutical Pvt. Ltd.

C-1, B-53, G.I.D.C. ESTATE,
NADIAD-387001 (GUJARAT) INDIA.
www.intracin.com

