

# K-GEN 10%

## Gentamicin Injection 100 mg/ml

### COMPOSITION :

#### Each ml contains :

Gentamicin Sulphate B.P	
eq. to Gentamicin Base	100 mg
Methyl Hydroxybenzoate B.P	0.18% w/v
(As Preservative)	
Propyl Hydroxybenzoate B.P	0.02% w/v
(As Preservative)	
Water for Injection B.P	q.s.

### INDICATION :

Gastrointestinal, respiratory infections, arthritis, poly-arthritis, meningitis, mastitis & metritis caused by Gentamicin sensitive bacteria, like E.Coli, Pasteurella, Klebsiella and salmonella spp., in Calves, Cattle, Horse, Goats and Sheep.

### DOSAGE AND ADMINISTRATION :

For Intramuscular administration:

General: Twice daily 1 ml per 20 - 40 kg body weight for 3 days.

### WITHDRAWAL PERIOD :

- For kidneys : 45 days.
- For meat : 7 days.
- For milk : 3 days.

### STORAGE :

Store at a temperature not exceeding 30°C.  
Protect from light.

IPPL

## Intracin

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,  
NADIAD-387001 (GUJARAT) INDIA.  
[www.intracin.com](http://www.intracin.com)

