

# SD-VET

## Sulfadimidine Injection B.P 33.3%

### COMPOSITION :

#### Each ml contains :

Sulfadimidine sodium B.P	333 mg
Benzyl alcohol B.P	1% v/v
Water for Injection B.P	q.s.

### INDICATION :

SD-VET is indicated for pneumonia, foot rot, diphtheria in Calves, actinobacillosis, coli-bacillosis, salmonellosis, pasteurellosis, enteritis, poli-arthritis, streptococcal mastitis, dysentery, lymphangitis, young's septicaemia and coccidiosis in Sheep and Goat, Cattle and Calves. For treatment of respiratory, lymphangitis, young's septicaemia and coccidiosis in Sheep and Goat, Cattle and Calves. For treatment of respiratory and coccidiosis infections in Poultry and Pigs.

### CONTRAINDICATION :

Contraindicated in known cases of hypersensitivity to sulphonamides and in animals with severe liver damage or blood dyscrasias.

### DOSAGE AND ADMINISTRATION :

For Subcutaneous and slow Intravenous administration.

**Cattle :** 3-6 ml/10 kg bodyweight the first day, followed by 3 ml per 10 kg. Body weight on following 2-5 days.

**Pigs :** 50 mg/kg bodyweight during 3-7 days.

**Poultry :** 100 mg/kg bodyweight during 3 days.

In cold weather the solution may deposit fine crystals. These will re-dissolved on gentle warming.

### WITHDRAWAL PERIOD :

Cattle, Sheep, Goat & Poultry : Meat : 18 days.

Cattle : Milk : 6.5 days.

Not for use in Sheep producing milk for human consumption.

Pigs : Meat : 42 days.

### STORAGE :

Store at a temperature not exceeding 30°C. Protect from light.

IPPL

**Intracin**

Pharmaceutical Pvt. Ltd.

C-1,B-53,G.I.D.C. ESTATE,  
NADIAD-387001 (GUJARAT) INDIA.  
www.intracin.com

